

Godman 1900. *Butleria pulchra*, *B. eryonas*, *B. lalage*, *B. octomaculata*, *B. faula*. Insecta.  
Lepidoptera-Rhopalocera. Vol II. [London: Published for the editors by R.H. Poter]. 1887-1901

5. ***Butleria pulchra***, sp. n. (Tab. XCII. figg. 7, 8, ♂.)

Alis fuscis, anticis maculis parvis quatuor discalibus (una infra cellulam triangularem), linea obliqua subapicali venis divisa, flavis; posticis maculis duabus conjunctis marginem externum versus, tertia minuta interiori, aurantiis: subtus fulvis, anticis maculis ut supra et margine interno late flavis, fascia lata a basi ad marginem externum extendente (maculis exceptis) fusca: posticis maculis paginae superioris fere obsoletis; ciliis posticarum aurantiis.

*Hab.* COSTA RICA, Irazu (*Rogers*).

This pretty species is also represented in our collection by a single male example, captured by Mr. Rogers on the slopes of the Volcano of Irazu. It approaches *B. agathocles* (Felder), from South America, in the disposition of the spots.

6. ***Butleria eryonas***. (Tab. XCII. figg. 9, 10, 11, ♂.)

*Cycloptides eryonas*, Hew. Ann. & Mag. Nat. Hist. (4) xx. p. 325 (1877) <sup>1</sup>.

*Butleria dolabella*, Plötz, Stett. ent. Zeit. 1884, p. 292 <sup>2</sup>.

Alis fuscis, anticis macula magna subquadrata inter ramos medianos primum et secundum, una minore in cellula triangularem, tertia margini externo propiore, punctisque duobus (superiore minuto) subapicalibus in linea transversa, hyalinis; posticis macula magna discali ovali ad cellulæ finem, flava: subtus fuscis, maculis omnibus paginae superioris multo majoribus; anticis ad apicem fascia transversa venis divisa, flava; posticis, costa ad basin, apice, et maculis tribus ad marginem externum flavis.

♀ nobis ignota.

*Hab.* PANAMA, Chiriqui <sup>2</sup> (*ex Staudinger, Ribbe* <sup>1</sup>), Veraguas (*Arcé*).

There are two specimens of this species in the Hewitson collection, both males, and we have three others of the same sex from the State of Panama, one of these latter having been sent us by Dr. Staudinger under the MS. name of *B. fimbriola*. For the genitalia of the male, see Tab. XCII. fig. 11.

7. ***Butleria lalage***, sp. n. (Tab. XCII. figg. 12, 13, ♂.)

Alis fuscis, anticis macula subtriangularem discali ad marginem externum dentata, altera obliqua minore subapicali et posticis plaga subovali discali, aurantiis: subtus aurantio-flavis, anticis plaga basali haud costam attingente, fascia lata curvata irregulari marginem externum versus, fuscis; ciliis posticarum flavis; subtus, palpis, thorace et abdomine pallide flavis.

*Hab.* MEXICO, Omilteme (*H. H. Smith*).

Mr. Smith procured us a single male example only of this species, from the neighbourhood of Omilteme, in Guerrero, from an altitude of 8000 feet. We possess an unnamed *Butleria* from Rio Janeiro which somewhat approaches it, but this insect has no subapical spot on the primaries.

8. ***Butleria octomaculata***, sp. n. (Tab. XCII. figg. 14, 15, ♂.)

Alis fuscis, anticis macula obliqua discali cellulam transeunte, altera minuta marginem propiore, tertia venis divisa subapicali, posticis subrotunda discali, omnibus pallide flavis: subtus fere ut supra, anticis macula discali literam V formante, posticis macula parva angulum analem versus, pallide flavis.

*Hab.* COSTA RICA, Irazu (*Rogers*).

We have but a single male example of *B. octomaculata*, sent us by our collector Mr. Rogers, from Irazu, where he captured it at an altitude of between 6000 and 7000 feet. It is allied to *B. cupavia*, Staud., from Bolivia.

9. **Butleria faula**, sp. n. (Tab. XCII. figg. 16, 17, ♂.)

Alis fuscis, macula subtriangulari discali ad marginem externum dentata, altera obliqua minore subapicali, et posticis macula elongata discali, secunda minuta infra eam et fere attingente, omnibus aurantiis: subtus aurantiis, anticis plaga basali haud ad costam extendente, fascia lata curvata irregulari marginem externum versus, fuscis; posticis obscure aurantiis, margine interno lato, fascia elongata per cellulam fasciolisque inter venas ad marginem externum, flavis; ciliis posticarum coloris ejusdem.

*Hab.* MEXICO, Guadalajara (*coll. Schaus*).

Mr. Schaus's collection contains a single male example of this species, from which the above description is taken. It is a close ally of *B. lalage*, but differs from it chiefly in having a long, narrow, discal band on the secondaries above and the underside of these wings otherwise coloured. *B. faula* is also very like *B. eryonas*, Hew., on the upperside, but differs considerably beneath.

B. Posterior tibiæ with a single pair of spurs.

PAMPHILA.

*Pamphila*, Fabricius, in Illiger's Mag. vi. p. 287 (1807); Watson, P. Z. S. 1893, p. 89.

*Steropes*, Boisduval, Voy. Astrolabe, Lép. p. 167 (1832).

*Carterocephalus*, Lederer, Verh. zool.-bot. Ges. Wien, ii. pp. 26, 49 (1853).